

**Original citation:**

Novak, L. A. and Gibbons, A. M. (1989) On perfect pairs of trees in a graph. University of Warwick. Department of Computer Science. (Department of Computer Science Research Report). (Unpublished) CS-RR-151

**Permanent WRAP url:**

<http://wrap.warwick.ac.uk/60845>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**A note on versions:**

The version presented in WRAP is the published version or, version of record, and may be cited as it appears here. For more information, please contact the WRAP Team at: [publications@warwick.ac.uk](mailto:publications@warwick.ac.uk)



<http://wrap.warwick.ac.uk/>

# \_\_\_\_Research report 151\_\_\_\_

## ON PERFECT PAIRS OF TREES IN A GRAPH

Ladislav Novak\*  
Alan Gibbons

(RR151)

Department of Computer Science  
University of Warwick  
Coventry CV4 7AL  
United Kingdom

\*Department of Electrical Engineering  
University of Novi Sad  
Yugoslavia

November 1989



# On Perfect Pairs of Trees in a Graph

Ladislav Novak

Department of Electrical Engineering, University of Novi Sad, Yugoslavia  
and

Alan Gibbons

Department of Computer Science, University of Warwick, England

## §1 Introduction

In many problems, especially those concerned with lumped circuits, pairs of (spanning) trees play an important role. Most "geometrical" properties of the set of all trees are related to the concept of distance between trees [1]. In that context several notions are introduced, in particular those of central tree [2], extremal tree [3] and maximally distant pairs of trees [3]. Some deep graph-theoretical concepts are related to the investigation of distances between trees, for example the concept of a principal partition of a graph [3] and that of topological degree of freedom [4]. Several important applications are also based on these notions [7]-[12].

Since Kishi and Kajitani realised the importance of introducing the concept of maximally distant trees [3] and Ohtsuki, Ishizaki and Watanabe promoted the concept of topological degree of freedom [4], many papers have been published which more closely characterise those properties of graphs connected with the space of trees [13]-[20]. The appearance of new concepts can be a spurious process and it may not at first be clear which are of value and whether the most useful definitions have been made. In this paper we introduce a new concept, that of a **perfect pair of trees** of a graph. We shall be particularly concerned with closely characterising the properties of this concept. We hope that our concept of a perfect pair of trees provides excellent intuitive insight within this area of study. Throughout this paper we shall be concerned only with 2-connected graphs.

## §2 Preliminaries

This section is devoted to some definitions and assertions related to material that follows. We presume that the reader is familiar with the following basic notions in graph theory : graph, edge, circuit and cutset. We take these to be primary notions that need not be defined. However we will define all other notions on the basis of these. Throughout we denote a graph by  $G$  and its edge set by  $E$ . The terms circuit, cutset, tree, cotree, forest and coforest will be used here to mean a subset

The following assertion plays a central role in this paper.

### Assertion 6

The following five statements are equivalent:

- i)  $t_2$  is maximally distant from  $t_1$
- ii) the fundamental circuit with respect to  $t_2$  defined by an edge in  $t_1^* \cap t_2^*$  contains no edges in  $t_1 \cap t_2$ .
- iii) the fundamental cutset with respect to  $t_1^*$  defined by an edge in  $t_1 \cap t_2$  contains no edges in  $t_1^* \cap t_2^*$ .
- iv)  $|t_1 \setminus t_2| = \text{rank } t_1^*$ .
- v) the number of edges in  $t_1 \cap t_2$  is equal to the maximal number of independent cutsets of the graph that belong entirely to the tree  $t_1$ .

### Proof

i)  $\Rightarrow$  ii)

Suppose that a fundamental circuit with respect to  $t_2$ , defined by an edge  $c \in t_1^* \cap t_2^*$ , contains an edge  $b \in t_1 \cap t_2$ , then  $t'_2 = (t_2 \setminus \{b\}) \cup \{c\}$  is a tree. Due to  $|t_1 \setminus t'_2| = |t_1 \setminus t_2| + 1$ , we conclude that the distance between  $t_1$  and  $t'_2$  is bigger than between  $t_1$  and  $t_2$ , which contradicts the assumption that  $t_2$  is maximally distant from  $t_1$ .

ii)  $\Leftrightarrow$  iii)

This is evident from the following well known statement: two edges belong to a circuit iff they both belong to the same cutset. Notice that in the present case, both circuit and cutset are in fact subsets of  $t_1 \setminus t_2$ .

ii)  $\Rightarrow$  iv)

If condition ii) is satisfied then fundamental circuits with respect to  $t_2$ , defined by edges in  $t_1^* \cap t_2^*$  contains only edges from  $t_2 \setminus t_1$ . Hence  $\text{rank } t_1^* \leq |t_2 \setminus t_1|$ . On the other hand, for any pair of trees  $(t_1, t_2)$   $\text{rank } t_1^* \geq |t_2 \setminus t_1|$ . (see Assertion 2, part i)). Therefore  $\text{rank } t_1^* = |t_2 \setminus t_1| = |t_1 \setminus t_2|$ .

iv)  $\Rightarrow$  i)

Suppose that the condition iv) is satisfied, and let  $t$  is an arbitrary tree. So, the following relations hold:

$$|t_1 \setminus t_2| = \text{rank } t_1^* \text{ (Condition iv))}$$

$$|t_1 \setminus t| \leq \text{rank } t_1^* \text{ (Assertion 2, part i))}$$

Consequently, for any tree  $t$ ,  $|t_1 \setminus t| \leq |t_1 \setminus t_2|$ .

(iv)  $\Leftrightarrow$  (v)

Let  $c$  be a number of vertices of the graph  $G$ . Also let  $p(t)$  be the number of connected components of the graph obtained by removing all edges from  $t$ , and let  $k(t)$  be the maximal number of independent cutsets in  $t_1$ . Then the following relations are always valid:

$$1) \text{ rank } t^* = c - p(t)$$

$$2) p(t) = k(t) + 1$$

$$3) c - 1 = |t|$$

$$4) |t| = |t \setminus t'| + |t \cap t'|, \text{ for 'arbitrary tree } t' \text{ in the graph.}$$

Applying these equations to the pair of trees  $(t_1, t_2)$  we immediately obtain the relation:

$$\text{rank } t_1^* = |t_1 \setminus t_2| + |t_1 \cap t_2| - k(t_1)$$

Obviously the relation  $\text{rank } t_1^* = |t_1 \setminus t_2|$  occurs iff  $|t_1 \cap t_2| = k(t_1)$  □

### Assertion 7

If  $t_2$  is maximally distant from  $t_1$ , then each common edge of  $t_1$  and  $t_2$  belongs to a cutset of the graph which is made of edges in  $t_2$  only.

### Proof

If  $t_2$  is maximally distant from  $t_1$  than according to Assertion 6, part(iii), an edge in  $t_1 \cap t_2$  can form a cutset only with the edges in  $t_1 \setminus t_2$ . But  $t_1 \cap t_2$  and  $t_1 \setminus t_2$  both belong to the tree  $t_2$  and consequently that cutset is made of the edges in  $t_2$  only. □

### Remark 1

The converse to Assertion 7 is generally not true. To see this, consider the graph shown in figure 3(a) and the two trees shown in (b) and (c) of the same figure. Common edges are marked as indicated in (d). It can be seen by inspection that all common edges of  $t_1$  and  $t_2$  belong to cutsets  $t_2$ , that is, the condition of Assertion 7 is fulfilled. Nevertheless, there exists a fundamental cutset (indicated by the curvy solid line in (c)) with respect to  $t_2$ , defined by a common edge of  $t_1$  and  $t_2$  which includes a common edge of the cotrees  $t_1^*$  and  $t_2^*$ . According to Assertion 6, part iii), we conclude that  $t_2$  is not maximally distant from  $t_1$ .




Figure 3

The following version of Assertion 7 is also true: If  $t_2$  is maximally distant from  $t_1$ , then each common edge of  $t_1^*$  and  $t_2^*$  belong to a circuit of the graph which is made of edges in  $t_1^*$  only.

#### §4 Perfect pairs of trees

A pair of trees  $(t_1, t_2)$  is defined to be a **perfect pair** of trees if both  $t_2$  is maximally distant from  $t_1$  and  $t_1$  is maximally distant from  $t_2$ .

As an immediate consequence of Assertion 6 we have the following theorem that gives several equivalent characterisations of perfect pairs.

##### Theorem 1

The following five statements are equivalent:

- i)  $(t_1, t_2)$  is a perfect pair
- ii) fundamental circuits with respect to  $t_1$  and  $t_2$  defined by edges in  $t_1^* \cap t_2^*$  contains no edges in  $t_1 \cap t_2$
- iii) fundamental cutsets with respect to  $t_1^*$  and  $t_2^*$  defined by edges in  $t_1 \cap t_2$  contains no edges in  $t_1^* \cap t_2^*$
- iv)  $\text{rank } t_1^* = |t_1 \setminus t_2| = |t_2 \setminus t_1| = \text{rank } t_2^*$
- v) the following three numbers, associated with the pair of trees  $(t_1, t_2)$  are equal:
  - the maximal number of independent cutsets of the graph that belong to  $t_1$
  - the maximal number of independent cutsets of the graph, that belong to  $t_2$
  - the number of common edges in  $t_1$  and  $t_2$ .

A pair  $(t_1, t_2)$  of trees is said to be a **maximally distant pair of trees** [3] if  $|t_1 \setminus t_2| \geq |t' \setminus t''|$  for every pair of trees  $(t', t'')$  in  $G$ .

### Assertion 8

A maximally distant pair is a perfect pair.

### Proof

If pair  $(t_1, t_2)$  is maximally distant then  $|t_1 \setminus t_2| \geq |t' \setminus t''|$  for every pair  $(t', t'')$  in the graph. For the particular cases:  $t'' = t_2$  and  $t' = t_1$ , we have  $|t_1 \setminus t_2| \geq |t_1 \setminus t''|$  for every  $t''$ , and  $|t_2 \setminus t_1| \geq |t_2 \setminus t'|$  for every  $t'$ .

Consequently, the pair  $(t_1, t_2)$  is a perfect pair.  $\square$

### Remark 2

The converse of Assertion 8 is not generally true. Figure 4(a) shows a perfect pair of trees with a corresponding distance of 5. That this pair is not a maximally distant pair is easily seen by noting that figure 4(b) shows a pair with a corresponding distance of 7. This second pair is obviously a maximally distant pair because the intersection of their edge sets is empty.




Figure 4

### Remark 3

We point out to the reader that the application of theorem 1 to the domain of maximally distant pairs (which is a restriction of the domain of perfect pairs, see remark 2) leads to a restriction of the theorem in the following sense: conditions ii)-v) are only necessary but not sufficient. More precisely we have the following corollary.

### Corollary (to Theorem 1 and Assertion 8)

If  $(t_1, t_2)$  is a maximally distant pair, then statements ii)-v) of theorem 1 hold.

A theorem related to maximally distant pairs similar to our corollary was presented in [21], (theorem 11.4). However, this theorem is only concerned with ii) and iii) of the corollary.

### Assertion 9

If  $t_2$  is maximally distant from  $t_1$  then each edge in  $t_2$  belongs to a fundamental circuit with respect to  $t_2$ , defined by an edge in  $t_1 \setminus t_2$ .

### Proof

Suppose that there exists an edge  $x \in t_2$ , that does not belong to a fundamental circuit with respect to



$t_2$  defined by edges in  $t_1 \setminus t_2$ . According to Assertion 4, such an edge as edge  $x$  can belong only to  $t_1 \cap t_2$ . On the other hand, in a 2-connected graph, each edge in  $t_2$  must belong to some fundamental circuit with respect to  $t_2^*$ . This means that  $x \in t_1 \cap t_2$  necessarily belongs to a fundamental circuit defined by an edge in  $t_1^* \setminus t_2^*$ , which contradicts the assumption that  $t_2$  is maximally distant from  $t_1$ .  $\square$

#### Remark 4

The converse statement to Assertion 9 is generally not true. To see this, consider the graph of figure 5(a) and the trees shown in (b) and (c) of that figure. Obviously  $\text{rank } t_1^* = 3 > \text{rank } t_2^* = 2 = |t_1 \setminus t_2|$ . So,  $t_2$  is not maximally distant from  $t_1$ . Nevertheless, each edge in  $t_2$  belongs to a fundamental circuit with respect to  $t_2$  defined by edges in  $t_1 \setminus t_2$ .




Figure 5

Denote by  $P$  the set of all distances between perfect pairs in the graph, and recall the definition of  $D$ .

#### Assertion 10

The following relation between  $P$  and  $D$  holds:

- (i)  $\min P \geq \min D$
- (ii)  $\max P = \max D$

#### Proof

(i) According to theorem 1, statement (iv), if  $t_1$  and  $t_2$  constitute a perfect pair, then they necessarily have the same diameter which is equal to their mutual distance apart. Hence,  $D \supseteq P$  and consequently  $\min P \geq \min D$ .

(ii) The distance between a pair of maximally distant pair is equal to  $\max D$ . But each maximally distant pair of trees is a perfect pair. Hence  $\max P = \max D$ .  $\square$

#### Remark 5

To prove the fact that strong inequality ( $\min P > \min D$ ) can occur, let us consider the graph of figure 6(a). Analysing the maximum number of independent cutsets that do not contain circuits, it is easy to see that the diameter of the graph is 4. There are exactly five trees with diameter 4 and these are shown figure 6(b) (each of them includes five independent cutsets of the graph). It can be checked immediately that the distance between any two of them is equal to 2. Hence, there is no perfect pair with distance 4.




Figure 6

## Conclusion

In this paper the new notion called a perfect pair of a graph was introduced. Several assertions were stated in order to closely characterise its properties. Also, several examples were included in order to help the reader gain intuitive insight.

## References

- [1] Hakimi, S L, "On trees of a graph and their generation", **J. Franklin Inst.**, vol 272, pp.347-359, November 1961.
- [2] Deo, N, "A central tree", **IEEE Trans.on Circuit Theory**, vol CT-13, pp.439-440, 1960.
- [3] Kishi, G and Kajitani, Y, "Maximally distant trees and principal partition of a linear graph", **IEEE Trans. on Circuit Theory**, vol.CT-16, pp.323-330, 1969.
- [4] Ohtsuki, T, Ishizaki, Y, and Watanabe, H, "Topological degrees of freedom and mixed analysis of electrical networks", **IEEE Trans. on Circuit Theory**, CT-17, pp.491-499, 1970.
- [5] Harary, F, Mokken, R J and Plantholt, M J, "Interpolation theorem for diameters of spanning trees", **IEEE Trans.on Circuits and Systems**, vol CAS-30, pp 429-431, 1983.
- [6] Novak, L A, "A few small theorems concerning pairs of matroid bases", to appear in **Int.J of Circuit Theory and Appl.** in 1990.
- [7] Lehman, A., "A solution of the Shannon switching game", **J. Soc. Indust. Appl. Math.**, vol.12, pp.687-724, 1964.
- [8] Bruno, J and Weinberg, L, "A constructive graph-theoretic solution of the shannon switching game", **IEEE Trans. on Circuits and Systems**, CAS-17, pp.74-81, 1970.
- [9] Chen, W K, and Chan, F N T, "On the unique solvability of linear active networks", **IEEE Trans. on Circuits and Systems**, pp.26-35, 1974.

- [10] Ozawa, T, "Topological conditions for the solvability of linear active networks", **Int. J. of Circuit Theory and Appl.**, pp.125-136, 1976.
- [11] Shinoda, S, "A minimization problem in network diagnosis", **Trans. IECEJ**, vol.53A, pp.569-570, 1970.
- [12] Tomizawa, N, "An algorithm for determining the maximally suspended pair of bases in a semi-monoid with applications to the diagnosis of networks", **Paper of Tech. Group on Circuit and System Theory of IECEJ**, CST74-80, pp.19-22, 1974.
- [13] Tomizawa, N, "An algorithm for determining the maximally distant pair of bases with matroidal constraints - generalized versions of Menger's theorem for matroids", **Papers of Tech. Group on Circuit and System Theory of IECEJ**, CST74-33, pp.17-24, 1974.
- [14] Kajitani, Y, and Kawamoto, T, "Graph theoretical degree of freedom in network analysis and all-central-tree graphs", **ibid**, CST76-94, pp.65-70, 1976.
- [15] Kawamoto, T, Kajitani, Y, and Shinoda, S, "New theorems on central trees described in connection with the principal partition of a graph", **ibid**, CST77-109, pp.63-68, 1977.
- [16] Kawamoto, T, and Shinoda, S, "Theorems on central trees in connection with the principal partition of a graph", **ibid**, CAS80-49, pp.41-45, 1980.
- [17] Kawamoto, T, Kajitani, Y, and Shinoda, S, "Eccentricity of the tree set", **Proc. of the IEEE Int. Symp. on Circuits and Systems**, paper R14-1, 1981.
- [18] Gao, S, and Chen, W K, "The hybrid method of network analysis and topological degree of freedom", **ibid**, pp.158-161, 1982.
- [19] Gao, S, and Chen, W K, "On graph classification with application", **Proc. 25th -Midwest Symp. on Circuits and Systems**, pp.244-248, 1982.
- [20] Lin, P M, "Complementary trees in circuit theory", **IEEE Trans. on Circuits and Systems**, CAS-27, pp.921-928, 1980.
- [21] Swamy, M N S and Thulasiraman, K, "Graphs, Networks and Algorithms", John Wiley and Sons, 1981.
- [22] Gibbons, A M, "Algorithmic Graph Theory", Cambridge University Press, 1985.




Figure 6

## Conclusion

In this paper the new notion called a perfect pair of a graph was introduced. Several assertions were stated in order to closely characterise its properties. Also, several examples were included in order to help the reader gain intuitive insight.

## References

- [1] Hakimi, S L, "On trees of a graph and their generation", **J. Franklin Inst.**, vol 272, pp.347-359, November 1961.
- [2] Deo, N, "A central tree", **IEEE Trans.on Circuit Theory**, vol CT-13, pp.439-440, 1960.
- [3] Kishi, G and Kajitani, Y, "Maximally distant trees and principal partition of a linear graph", **IEEE Trans. on Circuit Theory**, vol.CT-16, pp.323-330, 1969.
- [4] Ohtsuki, T, Ishizaki, Y, and Watanabe, H, "Topological degrees of freedom and mixed analysis of electrical networks", **IEEE Trans. on Circuit Theory**, CT-17, pp.491-499, 1970.
- [5] Harary, F, Mokken, R J and Plantholt, M J, "Interpolation theorem for diameters of spanning trees", **IEEE Trans.on Circuits and Systems**, vol CAS-30, pp 429-431, 1983.
- [6] Novak, L A, "A few small theorems concerning pairs of matroid bases", to appear in **Int.J of Circuit Theory and Appl.** in 1990.
- [7] Lehman, A., "A solution of the Shannon switching game", **J. Soc. Indust. Appl. Math.**, vol.12, pp.687-724, 1964.
- [8] Bruno, J and Weinberg, L, "A constructive graph-theoretic solution of the shannon switching game", **IEEE Trans. on Circuits and Systems**, CAS-17, pp.74-81, 1970.
- [9] Chen, W K, and Chan, F N T, "On the unique solvability of linear active networks", **IEEE Trans. on Circuits and Systems**, pp.26-35, 1974.

- [10] Ozawa, T, "Topological conditions for the solvability of linear active networks", **Int. J. of Circuit Theory and Appl.**, pp.125-136, 1976.
- [11] Shinoda, S, "A minimization problem in network diagnosis", **Trans. IECEJ**, vol.53A, pp.569-570, 1970.
- [12] Tomizawa, N, "An algorithm for determining the maximally suspended pair of bases in a semi-monoid with applications to the diagnosis of networks", **Paper of Tech. Group on Circuit and System Theory of IECEJ**, CST74-80, pp.19-22, 1974.
- [13] Tomizawa, N, "An algorithm for determining the maximally distant pair of bases with matroidal constraints - generalized versions of Menger's theorem for matroids", **Papers of Tech. Group on Circuit and System Theory of IECEJ**, CST74-33, pp.17-24, 1974.
- [14] Kajitani, Y, and Kawamoto, T, "Graph theoretical degree of freedom in network analysis and all-central-tree graphs", **ibid**, CST76-94, pp.65-70, 1976.
- [15] Kawamoto, T, Kajitani, Y, and Shinoda, S, "New theorems on central trees described in connection with the principal partition of a graph", **ibid**, CST77-109, pp.63-68, 1977.
- [16] Kawamoto, T, and Shinoda, S, "Theorems on central trees in connection with the principal partition of a graph", **ibid**, CAS80-49, pp.41-45, 1980.
- [17] Kawamoto, T, Kajitani, Y, and Shinoda, S, "Eccentricity of the tree set", **Proc. of the IEEE Int. Symp. on Circuits and Systems**, paper R14-1, 1981.
- [18] Gao, S, and Chen, W K, "The hybrid method of network analysis and topological degree of freedom", **ibid**, pp.158-161, 1982.
- [19] Gao, S, and Chen, W K, "On graph classification with application", **Proc. 25th -Midwest Symp. on Circuits and Systems**, pp.244-248, 1982.
- [20] Lin, P M, "Complementary trees in circuit theory", **IEEE Trans. on Circuits and Systems**, CAS-27, pp.921-928, 1980.
- [21] Swamy, M N S and Thulasiraman, K, "Graphs, Networks and Algorithms", John Wiley and Sons, 1981.
- [22] Gibbons, A M, "Algorithmic Graph Theory", Cambridge University Press, 1985.